

Operations on transcendental non-Archimedean metrics

Mingchen Xia

Dedicated to Bo Berndtsson.

ABSTRACT. We define the basic pluripotential-theoretic operations in terms of the transcendental theory of non-Archimedean metrics introduced in [DXZ23]. In particular, we establish that the analogue of Boucksom–Jonsson’s envelope conjecture holds in our theory.

CONTENTS

1. Introduction	1
2. Pluripotential theory on unibranch spaces	4
3. The theories of non-Archimedean metrics	7
4. Pluripotential-theoretic operations	13
5. Comparison with Boucksom–Jonsson’s theory	21
References	22

1. Introduction

Let X be a reduced irreducible unibranch compact Kähler space of dimension n and θ be a closed real $(1, 1)$ -form on X . Assume that $\text{PSH}(X, \theta) \neq \emptyset$. The purpose of this paper is to study the pluripotential theory of non-Archimedean metrics in the cohomology class of θ following the approach in [DXZ23].

1.1. Background. The non-Archimedean pluripotential theory has been developed by many authors for different purposes in the last three decades. The study of semipositive metrics on line bundles on non-Archimedean spaces dates back to the breakthrough of Zhang [Zha95]. In [CL06], Chambert-Loir initiated the study of the non-Archimedean Monge–Ampère measures. The pluripotential theory on non-Archimedean curves was developed in detail in the thesis of Thuillier

2020 *Mathematics Subject Classification.* Primary 32Q99.

The author benefited from discussions with Sébastien Boucksom. In particular, the possibility of obtaining Theorem 4.21 was suggested by him. The author would like to thank Tamás Darvas and Rémi Reboulet for their comments on the draft and the referee for numerous valuable suggestions. The author is supported by Knut och Alice Wallenbergs Stiftelse grant KAW 2021.0231.

[Thu05]. All these developments together lead to the natural question: to what extent is there a pluripotential theory on Berkovich spaces similar to its complex counterpart?

This question has drawn the attention of many authors. In particular, so far we know that there are two general approaches in the literature.

The first approach is the local approach, initiated by Chambert-Loir–Ducros [CLD12] and Gubler [Gub16]. In this approach, one defines the notions of differential forms and currents on a Berkovich space based on local tropicalizations. Then one can define the Monge–Ampère operator using the usual differential formula. The problem is that this approach only works for regular plurisubharmonic functions and completely fails when a singularity is presented. This is an essential obstacle for applications to complex geometry.

The second approach was initiated by Boucksom–Favre–Jonsson in [BFJ15, BFJ16], following the original ideas of Zhang. In the trivially valued case, this theory was further developed by Boucksom–Jonsson [BJ21]. In this approach, one first considers metrics obtained from semipositive models and then defines general global plurisubharmonic metrics using approximations. This approach allows singularities but does not work in the local situation. It does not yield a sheaf of plurisubharmonic metrics. Both approaches have their own drawbacks. We do not have a completely satisfactory non-Archimedean pluripotential theory yet.

1.2. The transcendental theory. The non-Archimedean pluripotential theory is closely related to complex geometry. We refer to [BBJ21] and the references therein. The basic idea is that the degeneration in the space of Kähler potentials can be characterized using global plurisubharmonic metrics on the non-Archimedean space. This idea has been intensively studied in [BBJ21, Li20, DX21, Xia23b].

We observe that there is a strong obstacle in applying the non-Archimedean geometry to complex geometry: the envelope conjecture [BJ21, Conjecture 5.14] of Boucksom–Jonsson remains open. This conjecture together with its various consequences is indispensable for various applications.

In our previous work [DXZ23], we introduced a different framework for the non-Archimedean global pluripotential theory based on the series of works starting from [BBJ21, RWN14]. Roughly speaking, we represent a non-Archimedean metric in $\text{PSH}^{\text{NA}}(X, \theta)$ by a *test curve* $\Gamma = (\Gamma_\tau)_\tau$, a concave curve of \mathcal{I} -model potentials. Due to the pathology caused by zero-mass, the rigorous definition requires a projective limit construction as well, we refer to Definition 3.5 for the rigorous definition. In this short introduction, we will only talk about test curves to simplify our presentation.

In the algebraic setting, when Boucksom–Jonsson’s theory is applicable, there is an obvious comparison map sending Γ to a function on the non-Archimedean space. See Definition 3.11 for the details. The main result in [DXZ23] asserts that when X is smooth, this comparison map is in fact a bijection to Boucksom–Jonsson’s theory, hence giving a complex representation of the non-Archimedean pluripotential theory. When X is only unibranch, Boucksom–Jonsson’s theory is still applicable. In this case, we prove the following result:

THEOREM 1.1 (=Corollary 5.3). *Assume that X is a reduced irreducible unibranch projective variety over \mathbb{C} and L is a pseudoeffective line bundle on X . Then the following are equivalent:*

- (1) *Boucksom–Jonsson’s envelope conjecture holds for (X, L) ;*
- (2) *The comparison map extends to a bijection from the space of non-Archimedean potentials in the sense of [DXZ23] to the corresponding space in the sense of [BJ21].*

Note that Boucksom–Jonsson’s envelope conjecture is known to hold when X is smooth, as confirmed in [BJ22]. A different proof was given in [DXZ23].

Using the comparison map, we can effectively translate the pluripotential-theoretic operations in Boucksom–Jonsson’s theory to our transcendental theory. It turns out that in most cases, these translations can be extended to more general situations where the comparison fails or when Boucksom–Jonsson’s theory is not applicable. This is exactly what we do in this paper. The final result is summarized in Table 1. It can be regarded as a duality between non-Archimedean geometry and convex geometry.

NA potentials	Test curves
Addition by a constant	Translation
Maximum	Concave envelope of maximum
Decreasing limit	Decreasing limit
Increasing limit	Increasing limit
Regularized supremum	Increasing limit of concave envelopes of maxima
Rescaling	Rescaling and reparametrization
Addition	Infimal involution
Restriction to subvarieties	Trace operator

TABLE 1. Correspondence non-Archimedean/convex

Most results from this table are quite straightforward to experts. The only exception is perhaps the last line (=Theorem 4.21), where we establish a quantitative version of the fact that a non-Archimedean plurisubharmonic metric is determined by its restriction to divisorial points. We hope that this result, after suitable extension, could serve to give a proof of Boucksom–Jonsson’s envelope conjecture in general.

As a particular case from this table, we have

THEOREM 1.2 (=Theorem 4.13). *The transcendental analogue of the envelope conjecture holds in the theory of [DXZ23].*

This result slightly generalizes [DXZ23, Theorem 1.2] in that the latter only concerns the regularized supremum of increasing nets while the former works for a general bounded from above family.

Given the fact that Boucksom–Jonsson’s envelope conjecture is widely open in general, we expect that our theory is more useful on normal varieties. In particular, we hope that the non-Archimedean theory of K-stability could be effectively translated to our theory. The very first step in this direction is carried out in [Xia23b]. On the other hand, our theory works more generally for transcendental classes on compact Kähler manifolds. It should be useful to the transcendental applications as well, for example in the study of cscK metrics as in [SD18] or probably in the study of the transcendental Morse inequality.

Most results from the current paper have been announced in several conferences. The author hopes that the details presented in the current paper are of interest to general readers as well. We assume that the readers are familiar with the complex pluripotential theory and Berkovich geometry.

1.3. Conventions. In the whole paper, the Monge–Ampère type products are always taken in the non-pluripolar sense. We adopt the convention that $dd^c = \frac{i}{2\pi}\partial\bar{\partial}$. A variety refers to a separated scheme of finite type over \mathbb{C} .

2. Pluripotential theory on unibranch spaces

Let X be a reduced irreducible unibranch compact Kähler space of dimension n and θ be a closed real $(1,1)$ -form on X . Here and in the sequel, a $(1,1)$ -form is always assumed to be smooth. Assume that $\text{PSH}(X, \theta) \neq \emptyset$. There is a well-developed pluripotential theory in this setting. For the Bedford–Taylor theory, we refer to [Dem85] and for the non-pluripolar theory, we refer to [Xia21a].

Note that we have assumed X to be irreducible. This is not a very severe restriction: since X is unibranch, its irreducible components are just its connected components. Every result in this paper extends in the obvious way to general reduced unibranch compact Kähler spaces.

2.1. Divisorial valuations.

Definition 2.1. A *prime divisor* over a reduced irreducible compact complex space Z is a connected smooth hypersurface $E \subseteq X'$, where $X' \rightarrow Z$ is a projective resolution.

Two prime divisors $E_1 \subseteq X'_1$ and $E_2 \subseteq X'_2$ over Z are *equivalent* if there is a common resolution $X'' \rightarrow X$ dominating both X'_1 and X'_2 such that the strict transforms of E_1 and E_2 coincide.

The set Z^{div} is the set of pairs (c, E) , where $c \in \mathbb{Q}_{>0}$ and E is an equivalence class of a prime divisor over Z . For simplicity, we will denote the pair (c, E) by $c \text{ord}_E$, although one should not really think of this object as a valuation unless Z is projective and irreducible.

Note that a prime divisor on Z does not always define a prime divisor over Z if Z is singular.

Definition 2.2. Given $\varphi \in \text{PSH}(X, \theta)$, we define its *associated non-Archimedean potential* as the map $\varphi^{\text{an}}: X^{\text{div}} \rightarrow \mathbb{R}$ given by

$$\varphi^{\text{an}}(c \text{ord}_E) := -c\nu(\varphi, E).$$

Here associated with E , there is a projective resolution $\pi: Y \rightarrow X$ such that E is a connected smooth hypersurface in Y . The notation $\nu(\varphi, E)$ means the generic Lelong number of $\pi^*\varphi$ along E . By Siu’s semicontinuity theorem, this quantity is independent of the choice of π .

2.2. The pluripotential theory.

Definition 2.3. Assume that X is smooth. Given $\varphi \in \text{PSH}(X, \theta)$, we define

$$P_\theta[\varphi] = \sup \left\{ \psi \in \text{PSH}(X, \theta) : \psi \leq 0, \int_X \theta_\varphi^n = \int_X \theta_\psi^n, \varphi \leq \psi + C \text{ for some } C \in \mathbb{R} \right\},$$

$$P_\theta[\varphi]_{\mathcal{I}} = \sup \{ \psi \in \text{PSH}(X, \theta) : \psi \leq 0, \mathcal{I}(k\varphi) = \mathcal{I}(k\psi) \text{ for all } k \in \mathbb{Z}_{>0} \}.$$

A potential $\varphi \in \text{PSH}(X, \theta)$ is *model* (resp. \mathcal{I} -*model*) if $\varphi = P_\theta[\varphi]$ (resp. $\varphi = P_\theta[\varphi]_{\mathcal{I}}$).

A potential $\varphi \in \text{PSH}(X, \theta)$ is \mathcal{I} -good if $\int_X \theta_\varphi^n > 0$ and $P_\theta[\varphi] = P_\theta[\varphi]_{\mathcal{I}}$.

Here $\mathcal{I}(k\varphi)$ denotes the multiplier ideal sheaf of $k\varphi$. For a detailed study of these notions, we refer to [DX21, DX22, Xia22].

Definition 2.4. We say a potential $\varphi \in \text{PSH}(X, \theta)$ is *model* (resp. \mathcal{I} -*model*, \mathcal{I} -*good*) if for a projective resolution $\pi: Y \rightarrow X$, $\pi^*\varphi \in \text{PSH}(Y, \pi^*\theta)$ is model (resp. \mathcal{I} -model, \mathcal{I} -good).

All three notions are independent of the choice of π .

Definition 2.5. Given $\varphi \in \text{PSH}(X, \theta)$, we define $P_\theta[\varphi], P_\theta[\varphi]_{\mathcal{I}} \in \text{PSH}(X, \theta)$ as follows: let $\pi: Y \rightarrow X$ be a projective resolution, then we require the following holds:

$$\pi^*P_\theta[\varphi] = P_{\pi^*\theta}[\pi^*\varphi], \quad \pi^*P_\theta[\varphi]_{\mathcal{I}} = P_{\pi^*\theta}[\pi^*\varphi]_{\mathcal{I}}.$$

By Zariski's main theorem, both $P_\theta[\varphi]$ and $P_\theta[\varphi]_{\mathcal{I}}$ are uniquely determined.

Both operations are clearly independent of the choice of π .

Definition 2.6. Two quasi-psh functions φ and ψ on X are P -*equivalent* if for some Kähler form ω on X such that $\varphi, \psi \in \text{PSH}(X, \omega)$ and both ω_φ and ω_ψ have positive masses, we have $P_\omega[\varphi] = P_\omega[\psi]$. We write $\varphi \sim_P \psi$ in this case.

Here and in the sequel, when we say a potential has positive mass, we always refer to the non-pluripolar mass.

Definition 2.7. Two quasi-psh functions φ and ψ on X are \mathcal{I} -*equivalent* if for some Kähler form ω on X such that both $\varphi, \psi \in \text{PSH}(X, \omega)$, we have $P_\omega[\varphi]_{\mathcal{I}} = P_\omega[\psi]_{\mathcal{I}}$. We write $\varphi \sim_{\mathcal{I}} \psi$ in this case.

Both conditions are independent of the choice of ω : for the \mathcal{I} -equivalence relation, this follows from [DX22, Theorem 2.13]. The statement for the P -equivalence relation is more tricky. As we do not need it in the paper, we omit the proof.

Proposition 2.8. *Suppose that $\varphi^i \in \text{PSH}(X, \theta)$ ($i \in I$) is a decreasing net of \mathcal{I} -model potentials and $\inf_i \varphi^i$ has positive mass. Then for all semipositive closed real $(1,1)$ -form ω on X , we have*

$$(2.1) \quad P_{\theta+\omega} \left[\inf_{i \in I} \varphi^i \right]_{\mathcal{I}} = \inf_{i \in I} P_{\theta+\omega}[\varphi^i]_{\mathcal{I}}.$$

The proof relies on the d_S -pseudometric introduced in [DDNL21].

PROOF. We may assume that X is smooth. In fact, we can always take a projection resolution of singularities $\pi: Y \rightarrow X$, then as explained in [Xia21a, Section 3.5], we have a canonical bijection $\text{PSH}(X, \theta) \cong \text{PSH}(Y, \pi^*\theta)$. Moreover, under this bijection, the pluripotential-theoretic operations on both spaces correspond to each other. So it suffices to prove (2.1) with Y , $\pi^*\theta$, $\pi^*\omega$ and $\pi^*\varphi^i$ in place of X , θ , ω and φ^i . This reduces the problem to the case where X is smooth. In the sequel, we will omit this kind of arguments.

Observe that both sides of (2.1) are \mathcal{I} -model in $\text{PSH}(X, \theta + \omega)$, so it suffices to show that they have the same generic Lelong numbers. Fix a prime divisor E

over X . As shown in [DDNL21, Proposition 4.8]¹, $P_{\theta+\omega}[\varphi^i]_{\mathcal{I}} \xrightarrow{d_S} \inf_{i \in I} P_{\theta+\omega}[\varphi^i]_{\mathcal{I}}$. Hence, by [Xia22, Theorem 0.10],

$$\lim_{i \in I} \nu(\varphi^i, E) = \nu \left(\inf_{i \in I} P_{\theta+\omega}[\varphi^i]_{\mathcal{I}}, E \right).$$

By [DDNL21, Proposition 4.8] again, $\varphi^i \xrightarrow{d_S} \inf_{i \in I} \varphi^i$. It follows from [Xia22, Theorem 0.10] that

$$\lim_{i \in I} \nu(\varphi^i, E) = \nu \left(\inf_{i \in I} \varphi^i, E \right).$$

We conclude (2.1). \square

Lemma 2.9. *Suppose that $\varphi^i \in \text{PSH}(X, \theta)$ ($i \in I$) is a non-empty family of \mathcal{I} -good potentials, uniformly bounded from above. Then $\sup^*_{i \in I} \varphi^i$ is also \mathcal{I} -good.*

In the proof below, $V_\theta := \sup\{\varphi \in \text{PSH}(X, \theta), \varphi \leq 0\}$.

PROOF. We may assume that X is smooth. Assume that this result is known when I is finite. By Choquet's lemma, we therefore assume that $I = \mathbb{Z}_{>0}$ and φ^i is an increasing sequence. In this case, the result follows from [Xia21b, Theorem 4.6].

It remains to handle the case where I has two elements. Let us take $\varphi, \psi \in \text{PSH}(X, \theta)$ be two \mathcal{I} -good potentials, we need to show that $\varphi \vee \psi$ is \mathcal{I} -good. By [DX21, Theorem 3.8], we may take sequences $\varphi_j, \psi_j \in \text{PSH}(X, \theta)$ converging to φ, ψ respectively with respect to d_S . It remains to argue that $\varphi_j \vee \psi_j \xrightarrow{d_S} \varphi \vee \psi$. This clearly follows from the following assertion: if $\varphi, \psi, \eta \in \text{PSH}(X, \theta)$, then

$$d_S(\varphi \vee \eta, \psi \vee \eta) \leq C_n d_S(\varphi, \psi)$$

for some constant $C_n > 0$ depending only on n . By [DDNL21, Proposition 3.5], we may assume that $\varphi \leq \psi$. Using the definition of d_S in [DDNL21], this would follow if we can show

$$d_1(\varphi \vee \eta \vee (V_\theta - C), \psi \vee \eta \vee (V_\theta - C)) \leq d_1(\varphi \vee (V_\theta - C), \psi \vee (V_\theta - C)),$$

which is just [Xia23a, Proposition 4.12] (Proposition 6.8 in the arXiv version). \square

Lemma 2.10. *Let ω be a semipositive closed real $(1, 1)$ -form on X . Let $\varphi^j \in \text{PSH}(X, \theta + \epsilon_j \omega)$ be a decreasing sequence of model potentials, where ϵ_j is a decreasing sequence converging to 0. Define $\varphi := \inf_j \varphi^j$.*

- (1) $\int_X \theta_\varphi^n = \lim_{j \rightarrow \infty} \int_X (\theta + \epsilon_j \omega + \text{dd}^c \varphi^j)^n$.
- (2) Assume that φ has positive mass. Then for any prime divisor E over X ,

$$\lim_{j \rightarrow \infty} \nu(\varphi^j, E) = \nu(\varphi, E).$$

PROOF. We may assume that X is smooth.

In this case, the result is proved in [DX24, Proposition 2.1]. \square

¹The result is stated for a sequence, but the proof works for a net as well.

2.3. The trace operator. The results in this section are simple extensions of the results established in [DX24]. The proofs are almost identical, so we omit the details.

Assume that X is smooth. Let E be an irreducible reduced closed subspace of X of dimension m .

Consider $\varphi \in \text{PSH}(X, \theta)$ with generic Lelong number $\nu(\varphi, E) = 0$. The trace operator gives a canonical and non-trivial way to restrict φ to E , different from the naive restriction $\varphi|_E$.

The problem with the naive restriction is that even under the assumption $\nu(\varphi, E) = 0$ it can happen that $\varphi|_E \equiv -\infty$. Moreover, the naive restriction is usually poorly behaved with respect to sequences of quasi-psh functions. The trace operator is a novel way of solving these issues.

Let $\varphi \in \text{PSH}(X, \theta)$ such that $\nu(\varphi, E) = 0$. We will define $\text{Tr}_E(\varphi)$ as a qpsf function on the normalization \tilde{E} of E , well-defined up to \mathcal{I} -equivalence. The definition is given in [DX24], which we recall now.

Definition 2.11. Suppose that φ has analytic singularities, then we let $\text{Tr}_E(\varphi) = \varphi|_{\tilde{E}}$. In general, fix a Kähler form ω on X and take a quasi-equisingular approximation $(\varphi_j)_j$ of φ in $\text{PSH}(X, \theta + \omega)$. We define $\text{Tr}_E(\varphi)$ as a d_S -limit of $\varphi_j|_{\tilde{E}}$.

Note that $\text{Tr}_E(\varphi)$ is not identical to $\text{Tr}_E^\theta(\varphi)$ in [DX24], but they are \mathcal{I} -equivalent when the latter has positive volume. Hence we can freely apply the results in [DX24].

The following result is proved in [DX24, Section 3].

Proposition 2.12. *Given $\varphi \in \text{PSH}(X, \theta)$ with $\nu(\varphi, E) = 0$, its trace operator $\text{Tr}_E(\varphi)$ as a quasi-psh function on \tilde{E} is well-defined up to \mathcal{I} -equivalence. Moreover, $\text{Tr}_E(\varphi)$ depends only on the \mathcal{I} -equivalence class of φ .*

Moreover, consider a decreasing sequence of \mathcal{I} -good potentials $\varphi_j \in \text{PSH}(X, \theta)$ with pointwise limit φ satisfying $\nu(\varphi, E) = 0$. Suppose that $\varphi_j \xrightarrow{d_S} \varphi$, then we have

$$\text{Tr}_E(\varphi_j) \xrightarrow{d_S} \text{Tr}_E(\varphi).$$

Here $\varphi_j \xrightarrow{d_S} \varphi$ means the same holds after pulling-back to an arbitrary projective resolution of singularities.

In general it is desirable to have a definition of $\text{Tr}_E(\varphi)$ when X is just unibranch. Unfortunately, due to the lack of Demailly regularization on X , the author cannot define the trace operator when E is contained in the singular locus of X . Subsequently, in Section 4.7, we have to assume that our Kähler space is smooth.

3. The theories of non-Archimedean metrics

In this section, we briefly recall two different theories of non-Archimedean metrics.

3.1. Boucksom–Jonsson theory. In this section, we recall the basic concept in Boucksom–Jonsson’s theory as in [BJ21].

Let X be an irreducible reduced variety over \mathbb{C} of dimension n . Let X^{an} denote the Berkovich analytification X^{an} of X with respect to the trivial valuation on \mathbb{C} .

The set of real valuations on $\mathbb{C}(X)$ trivial on \mathbb{C} is denoted by X^{val} . The center of a valuation v is the scheme-theoretic point $c = c(v)$ of X such that $v \geq 0$ on

$\mathcal{O}_{X,c}$ and $v > 0$ on the maximal ideal $\mathfrak{m}_{X,c}$ of $\mathcal{O}_{X,c}$. The center is unique if exists. It exists if X is proper.

In the remaining of this section, we assume that X is projective.

As a set, X^{an} is the set of semi-valuations on X , in other words, real-valued valuations v on irreducible reduced subvarieties Y in X that is trivial on \mathbb{C} . We call Y the *support* of the semi-valuation v . In other words,

$$X^{\text{an}} = \coprod_Y Y^{\text{val}}.$$

We will write $v_{\text{triv}} \in X^{\text{an}}$ for the trivial valuation on X : $v_{\text{triv}}(f) = 0$ for any $f \in \mathbb{C}(X)^\times$. See [Ber93] for more details.

We will be most interested in divisorial valuations. Recall that a divisorial valuation on X is a valuation of the form $c \text{ord}_E$, where $c \in \mathbb{Q}_{>0}$ and E is a prime divisor over X . The set of divisorial valuations on X is denoted by X^{div} . This notation is compatible with that in Section 2.1.

Given any coherent ideal \mathfrak{a} on X and any $v \in X^{\text{an}}$, we define

$$(3.1) \quad v(\mathfrak{a}) := \min\{v(f) : f \in \mathfrak{a}_{c(v)}\} \in [0, \infty],$$

where $c(v)$ is the center of the valuation v on X .

Given any valuation v on X , the Gauss extension of v is a valuation $\sigma(v)$ on $X \times \mathbb{A}^1$:

$$\sigma(v) \left(\sum_i f_i t^i \right) := \min_i (v(f_i) + i).$$

Here t is the standard coordinate on $\mathbb{A}^1 = \text{Spec } \mathbb{C}[t]$. The key property is that when v is a divisorial valuation, then so is $\sigma(v)$. See [BHJ17, Lemma 4.2].

Non-Archimedean plurisubharmonic functions. Let X be an irreducible complex projective variety of dimension n and L be a holomorphic pseudoeffective \mathbb{Q} -line bundle on X . Through the GAGA morphism $X^{\text{an}} \rightarrow X$ of ringed spaces, L can be pulled-back to an analytic line bundle L^{an} on X .

Following [BJ21, Definition 2.18], we define $\mathcal{H}_{\mathbb{Q}}^{\text{gf}}(L)$, the set of (*rational*) *generically finite Fubini-Study functions* $\phi : X^{\text{an}} \rightarrow [-\infty, \infty)$, that are of the following form:

$$(3.2) \quad \phi = \frac{1}{m} \max_j \{\log |s_j| + \lambda_j\}.$$

Here $m \in \mathbb{Z}_{>0}$ is an integer such that $L^{\otimes m}$ is a line bundle, the s_j 's are a finite collection of non-vanishing sections in $H^0(X, L^{\otimes m})$, and $\lambda_j \in \mathbb{Q}$. We followed the convention of Boucksom–Jonsson by writing $\log |s_j|(v) = -v(s_j)$.

Now we come to the main definition of this paragraph:

Definition 3.1 ([BJ21, Definition 4.1]). A psh metric on L^{an} is a function $\phi : X^{\text{an}} \rightarrow [-\infty, \infty)$ that is not identically $-\infty$, and is the pointwise limit of a decreasing net $(\phi_i)_{i \in I}$, where $\phi_i \in \mathcal{H}_{\mathbb{Q}}^{\text{gf}}(L_i^{\text{an}})$ for some \mathbb{Q} -line bundles L_i on X satisfying $c_1(L_i) \rightarrow c_1(L)$ in $\text{NS}^1(X)_{\mathbb{R}}$.

3.2. The transcendental theory. Let X be a reduced irreducible unibranch compact Kähler space of dimension n and θ be a closed real $(1,1)$ -form on X . Assume that $\text{PSH}(X, \theta)$ is non-empty.

We briefly recall the transcendental theory of non-Archimedean metrics introduced in [DXZ23].

Definition 3.2. The space $\text{PSH}^{\text{NA}'}(X, \theta)$ is defined as the set consisting of maps $\Gamma: (-\infty, \Gamma_{\max}) \rightarrow \text{PSH}(X, \theta)$ for some $\Gamma_{\max} \in \mathbb{R}$ satisfying the following conditions:

- (1) for each $\tau \in (-\infty, \Gamma_{\max})$, Γ_τ is \mathcal{I} -model;
- (2) the map $(-\infty, \Gamma_{\max}) \ni \tau \mapsto \Gamma_\tau$ is decreasing and concave.

We write

$$\Gamma_{-\infty} := \sup_{\tau \in (-\infty, \Gamma_{\max})}^* \Gamma_\tau.$$

We define $\text{PSH}^{\text{NA}}(X, \theta)_{>0}$ as the subset of $\text{PSH}^{\text{NA}'}(X, \theta)$ consisting of Γ satisfying furthermore

$$(3) \int_X \theta_{\Gamma_{-\infty}}^n > 0.$$

The curves $\Gamma \in \text{PSH}^{\text{NA}'}(X, \theta)$ are sometimes known as *test curves*. Here we choose the Greek letter Γ simply because it resembles the reflection of ℓ , the notation usually used to represent a geodesic ray, in view of their duality.

The object that we are eventually interested in is the space $\text{PSH}^{\text{NA}}(X, \theta)$ defined in Definition 3.5. The set $\text{PSH}^{\text{NA}'}(X, \theta)$ only plays an auxiliary role. In the notation $\text{PSH}^{\text{NA}}(X, \theta)_{>0}$, we did not put a prime in the index. Intuitively we would like to think of $\text{PSH}^{\text{NA}}(X, \theta)_{>0}$ as a subset of $\text{PSH}^{\text{NA}}(X, \theta)$ instead of a subset of $\text{PSH}^{\text{NA}'}(X, \theta)$. See Lemma 3.10.

Remark 3.3. Sometimes it is convenient to extend the domain of definition of $\Gamma \in \text{PSH}^{\text{NA}'}(X, \theta)$ to the whole \mathbb{R} as follows: for $\tau > \Gamma_{\max}$, we simply set $\Gamma_\tau = -\infty$ while for $\tau = \Gamma_{\max}$, we let

$$\Gamma_{\Gamma_{\max}} = \inf_{\tau < \Gamma_{\max}} \Gamma_\tau.$$

In particular, for each $x \in X$, the function $\mathbb{R} \ni \tau \mapsto \Gamma_\tau(x)$ is either constantly $-\infty$, or a proper usc concave function. We will frequently make use of this convention without further explanation.

Lemma 3.4. *Assume that $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$, then $\int_X \theta_{\Gamma_\tau}^n > 0$ for any $\tau \in (-\infty, \Gamma_{\max})$.*

PROOF. Fix $\tau \in (-\infty, \Gamma_{\max})$, we want to show that

$$(3.3) \quad \int_X \theta_{\Gamma_\tau}^n > 0.$$

We may assume that X is smooth.

By assumption, $\Gamma_{-\infty}$ has positive mass. By [DDNL18, Theorem 2.3], we have

$$\int_X \theta_{\Gamma_{-\infty}}^n = \lim_{\tau \rightarrow -\infty} \int_X \theta_{\Gamma_\tau}^n.$$

In particular, for a sufficiently small $\tau_0 < \tau$, we have

$$\int_X \theta_{\Gamma_{\tau_0}}^n > 0.$$

Now take $\tau' \in (\tau, \Gamma_{\max})$ and $t \in (0, 1)$ so that

$$\tau = (1-t)\tau' + t\tau_0.$$

From the concavity of Γ , we find that

$$\Gamma_\tau \geq (1-t)\Gamma_{\tau'} + t\Gamma_{\tau_0}.$$

By the monotonicity theorem [WN19],

$$\int_X \theta_{\Gamma_\tau}^n \geq \int_X \theta_{(1-t)\Gamma_{\tau'} + t\Gamma_{\tau_0}}^n \geq t^n \int_X \theta_{\Gamma_{\tau_0}}^n > 0$$

and (3.3) follows. \square

Definition 3.5. Given any Kähler form ω on X , we define transition maps

$$\begin{aligned} p_{\theta, \theta + \omega} &: \text{PSH}^{\text{NA}'}(X, \theta) \rightarrow \text{PSH}^{\text{NA}'}(X, \theta + \omega), \\ p_{\theta, \theta + \omega} &: \text{PSH}^{\text{NA}}(X, \theta)_{>0} \rightarrow \text{PSH}^{\text{NA}}(X, \theta + \omega)_{>0} \end{aligned}$$

as follows: the image of $\Gamma: (-\infty, \Gamma_{\max}) \rightarrow \text{PSH}(X, \theta)$ is given by

$$p_{\theta, \theta + \omega}(\Gamma): (-\infty, \Gamma_{\max}) \rightarrow \text{PSH}(X, \theta + \omega), \quad \tau \mapsto P_{\theta + \omega}[\Gamma_\tau]_{\mathcal{I}}.$$

Note that $\{\text{PSH}^{\text{NA}'}(X, \theta + \omega), p_{\theta, \theta + \omega}\}_\omega$ and $\{\text{PSH}^{\text{NA}}(X, \theta + \omega)_{>0}, p_{\theta, \theta + \omega}\}_\omega$ form projective systems, where the set of Kähler forms ω 's is ordered by the reverse of the usual ordering. We define

$$(3.4) \quad \text{PSH}^{\text{NA}}(X, \theta) := \varprojlim_{\omega} \text{PSH}^{\text{NA}'}(X, \theta + \omega),$$

where the projective limit is taken in the category of sets.

We will denote the components of an element $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ by $\Gamma^{\theta + \omega} \in \text{PSH}^{\text{NA}'}(X, \theta + \omega)$. Observe that $\Gamma_{\max}^{\theta + \omega}$ is independent of the choice of ω , we will denote this common value by Γ_{\max} .

Remark 3.6. The extension in Remark 3.3 is compatible with transition maps in the following sense: Suppose that $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$, then for any Kähler forms ω and ω' on X ,

$$P_{\theta + \omega + \omega'}[\Gamma_{\Gamma_{\max}^{\theta + \omega}}]_{\mathcal{I}} = \Gamma_{\Gamma_{\max}^{\theta + \omega + \omega'}}.$$

Lemma 3.7. *The natural maps*

$$\varprojlim_{\omega} \text{PSH}^{\text{NA}}(X, \theta + \omega)_{>0} \rightarrow \text{PSH}^{\text{NA}}(X, \theta) \rightarrow \varprojlim_{\omega} \text{PSH}^{\text{NA}}(X, \theta + \omega)$$

are both bijective.

In the sequel, we will be constantly making use of these identifications without further mentioning.

PROOF. It is clear that both maps are injective, so it suffices to verify that the map

$$\varprojlim_{\omega} \text{PSH}^{\text{NA}}(X, \theta + \omega)_{>0} \rightarrow \varprojlim_{\omega} \text{PSH}^{\text{NA}}(X, \theta + \omega)$$

is surjective. Consider an element $\Gamma \in \varprojlim_{\omega} \text{PSH}^{\text{NA}}(X, \theta + \omega)$ with components $\Gamma^{\theta + \omega} \in \text{PSH}^{\text{NA}}(X, \theta + \omega)$. For each ω , $\Gamma^{\theta + \omega}$ has components $\Gamma^{\theta + \omega, \theta + \omega + \omega'} \in \text{PSH}^{\text{NA}'}(X, \theta + \omega + \omega')$. It is easy to verify that Γ is the image of the following element in $\varprojlim_{\omega} \text{PSH}^{\text{NA}}(X, \theta + \omega)_{>0}$ with components $\Gamma^{\theta + \omega/2, \theta + \omega}$. \square

Also note that the transition maps give natural injections

$$(3.5) \quad \text{PSH}^{\text{NA}}(X, \theta) \hookrightarrow \text{PSH}^{\text{NA}}(X, \theta + \omega).$$

In particular, we can define

$$\text{QPSH}^{\text{NA}}(X) := \varinjlim_{\omega} \text{PSH}^{\text{NA}}(X, \theta + \omega),$$

where ω runs over all Kähler forms on X . This definition is clearly independent of the choice of θ .

Lemma 3.8. *The natural map*

$$(3.6) \quad \varinjlim_{\eta} \text{PSH}^{\text{NA}}(X, \theta + \eta)_{>0} \rightarrow \text{PSH}^{\text{NA}}(X, \theta)$$

is a bijection, where η runs over the set of semipositive closed real $(1, 1)$ -forms on X with positive total mass.

PROOF. The injectivity is trivial. In order to prove the surjectivity, let $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$. We want to show that Γ can be extended to an element in the domain of (3.6). For this purpose, take a semipositive closed real $(1, 1)$ -form η on X with positive total mass. Fix a Kähler form ω on X . We define

$$\Gamma_{\tau}^{\theta+\eta} := \inf_{k \in \mathbb{Z}_{>0}} \Gamma_{\tau}^{\theta+\eta+k^{-1}\omega}$$

for any $\tau \in (-\infty, \Gamma_{\max})$. By Lemma 2.10,

$$\int_X (\theta + \eta + \text{dd}^c \Gamma_{\tau}^{\theta+\eta})^n > 0$$

and

$$\nu(\Gamma_{\tau}^{\theta+\eta}, E) = \nu(\Gamma_{\tau}^{\theta+\eta+\omega}, E)$$

for any prime divisor E over X and any $\tau \in (-\infty, \Gamma_{\max})$. It is clear that $\Gamma^{\eta} \in \text{PSH}^{\text{NA}}(X, \theta + \eta)_{>0}$. The $(\Gamma^{\theta+\eta})_{\eta}$ defined in this way is clearly an element in the domain of (3.6) with image Γ . \square

Definition 3.9. The *non-pluripolar mass* of $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ is defined as the limit

$$\lim_{\omega} \int_X (\theta + \omega + \text{dd}^c \Gamma_{-\infty}^{\theta+\omega})^n,$$

where ω runs over the direct set of all Kähler forms on X . In other words, if we fix a Kähler form ω , the limit means

$$\lim_{k \rightarrow \infty} \int_X (\theta + k^{-1}\omega + \text{dd}^c \Gamma_{-\infty}^{\theta+k^{-1}\omega})^n,$$

Since the net is decreasing, the limit exists.

Lemma 3.10. *The image of the canonical injection*

$$\text{PSH}^{\text{NA}}(X, \theta)_{>0} \hookrightarrow \text{PSH}^{\text{NA}}(X, \theta)$$

is given by the set of $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ with positive non-pluripolar mass.

PROOF. It is clear that the image of an element in $\text{PSH}^{\text{NA}}(X, \theta)_{>0}$ has positive non-pluripolar mass. Conversely, take $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ with positive non-pluripolar mass. We want to construct $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ representing Γ .

Fix a Kähler form ω on X . Using the same arguments as Lemma 3.4, we find that

$$\lim_{k \rightarrow \infty} \int_X (\theta + k^{-1}\omega + \text{dd}^c \Gamma_{\tau}^{\theta+k^{-1}\omega})^n > 0$$

for any $\tau < \Gamma_{\max}$. We define

$$\Gamma'_\tau := \lim_{k \rightarrow \infty} \Gamma_\tau^{\theta+k^{-1}\omega}$$

for any $\tau < \Gamma_{\max}$. It follows from Lemma 2.10 that Γ' represents Γ . \square

3.3. Comparison between the two theories. Let X be a reduced irreducible unibranch projective variety over \mathbb{C} and L be a pseudoeffective line bundle on X . Recall that in this case, the Zariski unibranchness is equivalent to the analytic unibranchness, as proved in [Xia21a], so there is no ambiguity in the adjective *unibranch*.

Choose a smooth Hermitian metric h_0 on L and let $\theta = c_1(L, h_0)$. In this case, both the Boucksom–Jonsson theory and the transcendental theory make sense, so we can consider the problem of comparison.

3.3.1. The general case.

Definition 3.11. Given $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$, we define its *associated non-Archimedean potential* $\Gamma^{\text{an}}: X^{\text{div}} \rightarrow \mathbb{R}$ as follows:

$$(3.7) \quad \Gamma^{\text{an}}(c \text{ord}_E) := \sup_{\tau < \Gamma_{\max}} (\Gamma_\tau^{\theta+\omega, \text{an}}(c \text{ord}_E) + \tau)$$

for any Kähler form ω on X . Clearly, this map is independent of the choice of ω .

The map $\Gamma \mapsto \Gamma^{\text{an}}$ is compatible with the inclusions (3.5). In particular, given $\Gamma \in \text{QPSH}^{\text{NA}}(X)$, we can define Γ^{an} by choosing any representative of Γ .

It is clear that the map $\Gamma \mapsto \Gamma^{\text{an}}$ is injective.

3.3.2. On a smooth variety. In the sequel we assume that X is smooth. In this case, we have the techniques of multiplier ideal sheaves.

A potential $\varphi \in \text{PSH}(X, \theta)$ defines a potential $\varphi^{\text{an}} \in \text{PSH}(L^{\text{an}})$ as follows:

$$\varphi^{\text{an}}(v) := - \lim_{k \rightarrow \infty} \frac{1}{k} v(\mathcal{I}(k\varphi))$$

for any $v \in X^{\text{an}}$. As explained in [BBJ21, DXZ23], the limit exists and lies in $\text{PSH}(L^{\text{an}})$ and this function extends the non-Archimedean potential defined in Definition 2.2 on X^{div} .

We can then define a map

$$(3.8) \quad \text{an}: \text{PSH}^{\text{NA}}(X, \theta) \rightarrow \text{PSH}(L^{\text{an}})$$

as follows: take $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$, its image is given by

$$\Gamma^{\text{an}}: X^{\text{an}} \rightarrow [-\infty, \infty), \quad v \mapsto \sup_{\tau < \Gamma_{\max}} (\Gamma_\tau^{\theta+\omega, \text{an}}(v) + \tau)$$

for any Kähler form ω on X . Clearly, this map is independent of the choice of ω . The map Γ^{an} extends the corresponding definition in Definition 3.11.

THEOREM 3.12 ([DXZ23]). *Assume that X is smooth, then the map (3.8) is a bijection.*

Motivated by this bijection, we could translate the pluripotential-theoretic operations of $\text{PSH}(L^{\text{an}})$ to operations on the space $\text{PSH}^{\text{NA}}(X, \theta)$. Furthermore, in most cases, the resulting operations have natural extensions to the transcendental case as well. This is what we will carry out in the next section.

4. Pluripotential-theoretic operations

Let X be a reduced irreducible unibranch compact Kähler space of dimension n and θ be a closed real $(1,1)$ -form on X . Assume that $\text{PSH}(X, \theta) \neq \emptyset$. When necessary, we use θ' and θ'' to denote similar forms with the same properties as θ .

4.1. The partial order.

Definition 4.1. We define a partial order \leq on $\text{PSH}^{\text{NA}}(X, \theta)$ as follows: given $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$, we say $\Gamma \leq \Gamma'$ if for some (hence for all) Kähler form ω on X , we have $\Gamma_{\tau}^{\theta+\omega} \leq \Gamma'_{\tau}^{\theta+\omega}$ for all $\tau < \Gamma_{\max}$.

It is trivial to verify that this is indeed a partial order. Note that this partial order is compatible with the inclusions (3.5), so it defines a partial order on $\text{QPSH}^{\text{NA}}(X)$.

Lemma 4.2. *Suppose that $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$, then the following are equivalent:*

- (1) $\Gamma \leq \Gamma'$;
- (2) $\Gamma^{\text{an}}(\text{cord}_E) \leq \Gamma'^{\text{an}}(\text{cord}_E)$ for all $\text{cord}_E \in X^{\text{div}}$.

The proof is almost identical to [DXZ23, Theorem 3.12], we omit the details.

4.2. Addition by a constant.

Definition 4.3. Given $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ and $C \in \mathbb{R}$, we define the addition $\Gamma + C \in \text{PSH}^{\text{NA}}(X, \theta)$ as the element with components $(\Gamma + C)^{\theta+\omega} : (-\infty, \Gamma_{\max} + C) \rightarrow \text{PSH}(X, \theta + \omega)$ given by

$$(\Gamma + C)_{\tau}^{\theta+\omega} := \Gamma_{\tau-C}^{\theta+\omega}.$$

The associated non-Archimedean potential behaves as expected:

Proposition 4.4. *Given $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ and $C \in \mathbb{R}$, we have*

$$(4.1) \quad (\Gamma + C)^{\text{an}}(\text{cord}_E) = \Gamma^{\text{an}}(\text{cord}_E) + C$$

for all $\text{cord}_E \in X^{\text{div}}$ and

$$(\Gamma + C)_{\max} = \Gamma_{\max} + C.$$

This follows trivially from (3.7). We will omit similar proofs in the sequel.

Of course, this operation has the obvious compatibility in itself: given another constant $C' \in \mathbb{R}$, we have

$$(\Gamma + C) + C' = \Gamma + (C + C').$$

It is compatible with the partial order in the following sense: given another $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$, the following are equivalent:

- (1) $\Gamma \leq \Gamma'$;
- (2) $\Gamma + C \leq \Gamma' + C$.

4.3. Maximum.

Definition 4.5. Suppose that $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$, we define their *maximum* $\Gamma \vee \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ as the map $(-\infty, \Gamma_{\max} \vee \Gamma'_{\max}) \rightarrow \text{PSH}(X, \theta)$:

$$\tau \mapsto P_{\theta}[\text{EC}(\tau' \mapsto \Gamma_{\tau'} \vee \Gamma'_{\tau'})_{\tau}]_{\mathcal{I}},$$

where EC denotes the concave hull.

We need to verify that $\Gamma \vee \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$. The only non-trivial part is to show that the concave envelope $\text{EC}(\tau' \mapsto \Gamma_{\tau'} \vee \Gamma'_{\tau'})$ is a curve of θ -psh functions. For this purpose, we may assume that X is smooth and connected. According to the Ross–Witt Nyström correspondence ([**RWN14**, Section 5, Section 6] and [**DXZ23**, Theorem 2.6]), Γ and Γ' correspond to geodesic rays ℓ and ℓ' in $\text{PSH}(X, \theta)$: for any $t \geq 0$, we have

$$\ell_t = \sup_{\tau \in \mathbb{R}} (\Gamma_{\tau} + t\tau), \quad \ell'_t = \sup_{\tau \in \mathbb{R}} (\Gamma'_{\tau} + t\tau).$$

It follows that

$$\ell_t \vee \ell'_t = \sup_{\tau \in \mathbb{R}} (\Gamma_{\tau} \vee \Gamma'_{\tau} + t\tau).$$

Hence,

$$\text{EC}(\tau \mapsto \Gamma_{\tau} \vee \Gamma'_{\tau})_{\tau'} = \inf_{t \geq 0} (\ell_t \vee \ell'_t - t\tau')$$

and $\text{EC}(\tau \mapsto \Gamma_{\tau} \vee \Gamma'_{\tau})_{\tau'} \in \text{PSH}(X, \theta) \cup \{-\infty\}$ by Kiselman's principle.

By a similar argument, we have

Corollary 4.6. *Suppose that $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(\theta)_{>0}$. For any $c \text{ord}_E \in X^{\text{div}}$, we have*

$$-(\Gamma \vee \Gamma')_{\tau}^{\text{an}}(c \text{ord}_E) = \text{EC}(\tau' \mapsto (-\Gamma_{\tau'}^{\text{an}}(c \text{ord}_E)) \vee (-\Gamma'_{\tau'}^{\text{an}}(c \text{ord}_E)))_{\tau}.$$

In particular,

$$(4.2) \quad (\Gamma \vee \Gamma')^{\text{an}}(c \text{ord}_E) = \Gamma^{\text{an}}(c \text{ord}_E) \vee \Gamma'^{\text{an}}(c \text{ord}_E).$$

It follows from Corollary 4.6 and Lemma 4.2 that the maximum operation is compatible with the projective system (3.4). Hence, we can make the following definition in general:

Definition 4.7. Suppose that $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$, we define their *maximum* $\Gamma \vee \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$ as the element with components

$$(\Gamma \vee \Gamma')^{\theta+\omega} = \Gamma^{\theta+\omega} \vee \Gamma'^{\theta+\omega}.$$

The associated non-Archimedean potential behaves as expected:

Corollary 4.8. *Suppose that $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(\theta)$. For any $c \text{ord}_E \in X^{\text{div}}$, we have*

$$(4.3) \quad (\Gamma \vee \Gamma')^{\text{an}}(c \text{ord}_E) = \Gamma^{\text{an}}(c \text{ord}_E) \vee \Gamma'^{\text{an}}(c \text{ord}_E)$$

and

$$(\Gamma \vee \Gamma')_{\max} = \Gamma_{\max} \vee \Gamma'_{\max}.$$

The maximum operation is commutative and associative. In particular, finite (non-empty) maximum makes sense in the obvious way.

The maximum operation is compatible with the partial order in the following sense: given $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(\theta)$, the following are equivalent

- (1) $\Gamma \vee \Gamma' = \Gamma'$;
- (2) $\Gamma \leq \Gamma'$.

The maximum operation is compatible with the addition by a constant: if $C \in \mathbb{R}$, then

$$(\Gamma \vee \Gamma') + C = (\Gamma + C) \vee (\Gamma' + C).$$

4.4. Decreasing limit along a net.

Definition 4.9. Let $(\Gamma_i)_{i \in I}$ be a decreasing net in $\text{PSH}^{\text{NA}}(X, \theta)$. Assume that $\inf_{i \in I} \Gamma_{i, \max} > -\infty$. We define $\inf_{i \in I} \Gamma_i \in \text{PSH}^{\text{NA}}(X, \theta)$ as the element with components

$$\left(\inf_{i \in I} \Gamma_i \right)_{\tau}^{\theta + \omega} = \inf_{i \in I} \Gamma_{i, \tau}^{\theta + \omega}.$$

This gives an element in $\text{PSH}^{\text{NA}}(X, \theta)$ by Proposition 2.8 together with [DX22, Lemma 2.21(i)]².

When $\inf_{i \in I} \Gamma_{i, \max} = -\infty$, we could still formally define $\inf_{i \in I} \Gamma_i$ as a symbol $-\infty$. As one could easily verify, all of our pluripotential-theoretic operations admit natural extensions to $-\infty$.

Using [DXZ23, Lemma 3.13], we have

Lemma 4.10. *Let $(\Gamma_i)_{i \in I}$ be a decreasing net in $\text{PSH}^{\text{NA}}(X, \theta)$ with $\inf_{i \in I} \Gamma_i \in \text{PSH}^{\text{NA}}(X, \theta)$. Then for any $\text{cord}_E \in X^{\text{div}}$,*

$$(4.4) \quad \left(\inf_{i \in I} \Gamma_i \right)_{\text{max}}^{\text{an}}(\text{cord}_E) = \inf_{i \in I} \Gamma_i^{\text{an}}(\text{cord}_E).$$

Moreover,

$$\left(\inf_{i \in I} \Gamma_i \right)_{\text{max}} = \inf_{i \in I} \Gamma_{i, \text{max}}.$$

This operation is compatible with itself: if I and J are two non-empty directed set and Γ_{ij} is a decreasing net in $\text{PSH}^{\text{NA}}(X, \theta)$ indexed by $I \times J$ with $\inf_{i, j} \Gamma_{ij} \in \text{PSH}^{\text{NA}}(X, \theta)$, then

$$\inf_{i, j} \Gamma_{ij} = \inf_{i \in I} \inf_{j \in J} \Gamma_{ij} = \inf_{j \in J} \inf_{i \in I} \Gamma_{ij}.$$

This operation is compatible with the partial order in the following senses: if $\Gamma_i \in \text{PSH}^{\text{NA}}(X, \theta)$ ($i \in I$) is a decreasing net with $\inf_{i \in I} \Gamma_i \in \text{PSH}^{\text{NA}}(X, \theta)$. Then $\inf_{j \in I} \Gamma_j \leq \Gamma_i$ for all $i \in I$ and it is the biggest element in $\text{PSH}^{\text{NA}}(X, \theta)$ with this property. On the other hand, if $\Gamma'_i \in \text{PSH}^{\text{NA}}(X, \theta)$ ($i \in I$) is another decreasing net with $\inf_{i \in I} \Gamma'_i \in \text{PSH}^{\text{NA}}(X, \theta)$ such that $\Gamma_i \leq \Gamma'_i$ for all $i \in I$, then

$$\inf_{i \in I} \Gamma_i \leq \inf_{i \in I} \Gamma'_i.$$

This operation is compatible with the addition by a constant: if $C \in \mathbb{R}$, then

$$\inf_{i \in I} (\Gamma_i + C) = \left(\inf_{i \in I} \Gamma_i \right) + C.$$

4.5. Regularized supremum.

Definition 4.11. Let $(\Gamma^i)_{i \in I}$ be an increasing net in $\text{PSH}^{\text{NA}}(X, \theta)_{>0}$. We assume that $\sup_{i \in I} \Gamma_{\max}^i < \infty$. We define a map $\sup_{i \in I} \Gamma_{\max}^i : (-\infty, \sup_{i \in I} \Gamma_{\max}^i) \rightarrow \text{PSH}(X, \theta)$ as follows:

$$\left(\sup_{i \in I}^* \Gamma^i \right)_{\tau} = P_{\theta} \left[\sup_{i \in I}^* \Gamma_{\tau}^i \right].$$

By Lemma 2.9, it is easy to see that $\sup_{i \in I}^* \Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$.

²The result is stated for a sequence, but the proof extends to a net.

More generally, suppose that $\Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta)$ ($i \in I$) is an increasing net with $\sup_i \Gamma_{\max}^i < \infty$. We define $\sup^* \Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta)$ as the element with components

$$(4.5) \quad \left(\sup_{i \in I}^* \Gamma^i \right)^{\theta + \omega} = \sup_{i \in I}^* \Gamma^{i, \theta + \omega}$$

for any Kähler form ω on X . The compatibility follows from [DX21, Lemma 3.14].

Definition 4.12. Let $(\Gamma^i)_{i \in I}$ be a (non-empty) family in $\text{PSH}^{\text{NA}}(X, \theta)$ such that $\sup_{i \in I} \Gamma_{\max}^i < \infty$. Let \mathcal{J} denote the directed set of finite non-empty subsets of I ordered by inclusion. Then we define

$$\sup_{i \in I}^* \Gamma^i := \sup_{J \in \mathcal{J}}^* \left(\max_{j \in J} \Gamma^j \right).$$

Recall that the maximum is defined right after Corollary 4.8.

THEOREM 4.13. *Let $(\Gamma^i)_{i \in I}$ be a (non-empty) family in $\text{PSH}^{\text{NA}}(X, \theta)$ with $\sup_{i \in I} \Gamma_{\max}^i < \infty$. Then for any $\text{cord}_E \in X^{\text{div}}$, we have*

$$\left(\sup_{i \in I}^* \Gamma^i \right)^{\text{an}} (\text{cord}_E) = \sup_{i \in I} \Gamma^{i, \text{an}} (\text{cord}_E).$$

Moreover,

$$\left(\sup_{i \in I}^* \Gamma^i \right)_{\max} = \sup_{i \in I} \Gamma_{\max}^i.$$

This result is the transcendental analogue of Boucksom–Jonsson’s envelope conjecture. As explained in [DXZ23, Theorem 3.4], when X is smooth and projective, this result implies Boucksom–Jonsson’s envelope conjecture.

PROOF. By Corollary 4.8, we may assume that $(\Gamma^i)_{i \in I}$ is an increasing net. Furthermore, it is easy to reduce to the case where $\Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$. In fact, by definition,

$$\left(\sup_{i \in I}^* \Gamma^i \right)^{\text{an}} = \left(\sup_{i \in I}^* \Gamma^{i, \theta + \omega} \right)^{\text{an}}$$

for any Kähler form ω on X . A similar equation holds for $(\bullet)_{\max}$ as well. So we could replace Γ^i by $\Gamma^{i, \theta + \omega}$ and assume that $\Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$.

In this case, the result follows from [DXZ23, Lemma 3.13]. \square

In particular, this operation extends the finite maximum.

This operation is compatible with itself: if $(\Gamma^{i,j})_{i \in I, j \in J}$ is a (non-empty) family in $\text{PSH}^{\text{NA}}(X, \theta)$ with $\sup_{i \in I, j \in J} \Gamma_{\max}^{i,j} < \infty$. Then

$$\sup_{i \in I}^* \sup_{j \in J}^* \Gamma^{i,j} = \sup_{j \in J}^* \sup_{i \in I}^* \Gamma^{i,j}.$$

It is compatible with the partial order: Suppose that $(\Gamma^i)_{i \in I}, (\Gamma'^i)_{i \in I}$ are two families in $\text{PSH}^{\text{NA}}(X, \theta)$ with $\sup_{i \in I} \Gamma_{\max}^i < \infty$ and $\sup_{i \in I} \Gamma'_{\max}{}^i < \infty$ and $\Gamma^i \leq \Gamma'^i$ for each $i \in I$, then

$$\sup_{i \in I}^* \Gamma^i \leq \sup_{i \in I}^* \Gamma'^i.$$

It is compatible with addition by a constant: if $C \in \mathbb{R}$, then

$$\sup_{i \in I}^* (\Gamma^i + C) = \left(\sup_{i \in I}^* \Gamma^i \right) + C.$$

There is no non-trivial compatibility with the decreasing limits, as expected.

4.6. Rescaling and addition.

Definition 4.14. Let $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ and $\lambda \in \mathbb{R}_{>0}$, we define the rescaling $\lambda\Gamma \in \text{PSH}^{\text{NA}}(X, \lambda\theta)$ as the element with components $(\lambda\Gamma)^{\lambda\theta+\omega} : (-\infty, \lambda\Gamma_{\max}) \rightarrow \text{PSH}(X, \lambda\theta + \omega)$:

$$(\lambda\Gamma)_{\tau}^{\lambda\theta+\omega} := \lambda\Gamma_{\lambda^{-1}\tau}^{\theta+\lambda^{-1}\omega}$$

for any Kähler form ω on X .

Definition 4.15. Let $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ and $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')_{>0}$. We define $\Gamma + \Gamma' : (-\infty, \Gamma_{\max} + \Gamma'_{\max}) \rightarrow \text{PSH}(X, \theta + \theta')$ as

$$(\Gamma + \Gamma')_{\tau} := P_{\theta+\theta'} \left[\sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-t}) \right].$$

Observe that $\tau \mapsto \sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-t})$ is concave. In fact, this curve is just the concave involution of Γ and Γ' . From the general properties of infimal involution, we know that this map is concave and usc. For each fixed $x \in X$, $\tau \mapsto \sup_{t \in \mathbb{R}} (\Gamma_t(x) + \Gamma'_{\tau-t}(x))$ is either constantly $-\infty$ or proper. In particular, by [Roc70, Theorem 16.4] and Legendre–Fenchel duality, $\tau \mapsto \sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-t})$ is the same as the inverse Legendre transform of $\ell + \ell'$, where ℓ and ℓ' are the geodesic rays defined by Γ and Γ' . In particular, $\sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-t})$ is either in $\text{PSH}(X, \theta)$ or constantly $-\infty$ by Kiselman’s principle. By Lemma 2.9 and [Xia22, Proposition 2.8], we find that $\Gamma + \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$.

This argument also shows that for any $c \text{ord}_E \in X^{\text{div}}$, we have

$$(4.6) \quad (\Gamma + \Gamma')^{\text{an}}(c \text{ord}_E) = \Gamma^{\text{an}}(c \text{ord}_E) + \Gamma'^{\text{an}}(c \text{ord}_E).$$

Definition 4.16. Let $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ and $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')$. We define $\Gamma + \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta + \theta')$ as the element with components

$$(\Gamma + \Gamma')^{\theta+\omega} = \Gamma^{\theta+\omega} + \Gamma'^{\theta+\omega}.$$

Note that $\Gamma + \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta + \theta')$ by (4.6).

Lemma 4.17. *Suppose that $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$, $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')$, $\lambda \in \mathbb{R}_{>0}$. Then for any $c \text{ord}_E \in X^{\text{div}}$, we have*

$$(\lambda\Gamma)^{\text{an}}(c \text{ord}_E) = \lambda\Gamma^{\text{an}}(c \text{ord}_E), \quad (\Gamma + \Gamma')^{\text{an}}(c \text{ord}_E) = \Gamma^{\text{an}}(c \text{ord}_E) + \Gamma'^{\text{an}}(c \text{ord}_E).$$

Moreover,

$$(\lambda\Gamma)_{\max} = \lambda\Gamma_{\max}, \quad (\Gamma + \Gamma')_{\max} = \Gamma_{\max} + \Gamma'_{\max}.$$

Both operations are compatible with themselves: if $\lambda, \lambda' \in \mathbb{R}_{>0}$ and $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$, $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')$ and $\Gamma'' \in \text{PSH}^{\text{NA}}(X, \theta'')$, then

$$\lambda(\lambda'\Gamma) = (\lambda\lambda')\Gamma$$

and

$$(\Gamma + \Gamma') + \Gamma'' = \Gamma + (\Gamma' + \Gamma'').$$

The addition is clear commutative. Moreover, these two operations are compatible with each other:

$$\lambda(\Gamma + \Gamma') = \lambda\Gamma + \lambda\Gamma', \quad (\lambda + \lambda')\Gamma = \lambda\Gamma + \lambda'\Gamma.$$

Both operations are compatible with the partial order: if $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$, $\lambda \in \mathbb{R}_{>0}$, then $\Gamma \leq \Gamma'$ if and only if $\lambda\Gamma \leq \lambda\Gamma'$. Suppose that $\Gamma^1, \Gamma^2 \in \text{PSH}^{\text{NA}}(X, \theta)$, $\Gamma'^1, \Gamma'^2 \in \text{PSH}^{\text{NA}}(X, \theta')$ and $\Gamma^1 \leq \Gamma^2$, $\Gamma'^1 \leq \Gamma'^2$, then $\Gamma^1 + \Gamma'^1 \leq \Gamma^2 + \Gamma'^2$.

Both operations are compatible with addition by a constant: if $C \in \mathbb{R}$, $\lambda \in \mathbb{R}_{>0}$, $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ and $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')$, then

$$\lambda(\Gamma + C) = \lambda\Gamma + \lambda C$$

and

$$(\Gamma + \Gamma') + C = \Gamma + (\Gamma' + C).$$

Both operations satisfy obvious inequalities with respect to the limit operations. We leave the details to the readers.

4.7. The trace operator. We assume in addition that X is smooth. Recall that in this case, we have a natural bijection (3.8). Fix an irreducible reduced closed analytic subspace E of X .

Definition 4.18. Assume that X is smooth. We say $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ has *non-trivial restriction* to E if there is a small enough $C \in \mathbb{R}$ so that $\nu(\Gamma_C^{\theta+\omega}, E) = 0$ for any Kähler form ω on X .

This definition is motivated by the following observation.

Lemma 4.19. *Assume that X is smooth and projective. Let $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$. Then the following are equivalent:*

- (1) $\Gamma^{\text{an}}(v_{E, \text{triv}}) \neq -\infty$;
- (2) $\Gamma^{\text{an}}|_{E^{\text{an}}} \neq -\infty$;
- (3) Γ has non-trivial restriction to E .

Here $v_{E, \text{triv}}$ denotes the trivial valuation of $\mathbb{C}(E)$.

PROOF. The equivalence between (1) and (2) is a simple consequence of the maximum principle [BJ21, Lemma 1.4(i)].

To see the equivalence between (1) and (3), it suffices to observe that for any $\varphi \in \text{PSH}(X, \theta)$,

$$\varphi^{\text{an}}(v_{E, \text{triv}}) = \begin{cases} -\infty, & \text{if } \nu(\varphi, E) > 0; \\ 0, & \text{if } \nu(\varphi, E) = 0. \end{cases}$$

□

Definition 4.20. Assume that X is smooth. Assume that Γ has non-trivial restriction to E , we define the *trace operator* of Γ along E as the element $\text{Tr}_E(\Gamma) \in \text{PSH}^{\text{NA}}(\tilde{E}, \theta|_{\tilde{E}})$ defined as follows: for any Kähler form ω' on Y and any Kähler form ω on X satisfying $\omega' \geq \omega|_{\tilde{Y}}$, the component $\text{Tr}_E(\Gamma)^{\theta|_{\tilde{E}}+\omega'} \in \text{PSH}^{\text{NA}}(\tilde{E}, \theta|_{\tilde{E}} + \omega')$ is given by

$$\tau \mapsto P_{\theta|_{\tilde{E}}+\omega'}[\text{Tr}_E(\Gamma_\tau^{\theta+\omega})],$$

where $\tau \in (-\infty, (\text{Tr}_E(\Gamma))_{\max})$ and $(\text{Tr}_E(\Gamma))_{\max} = \sup\{\tau \in \mathbb{R} : \nu(\Gamma_\tau^{\theta+\omega'}, E) = 0\}$ for any Kähler form ω'' on X .

We remind the readers that \tilde{E} is the normalization of E . It is a normal Kähler space by [GK20, Proposition 3.5].

The non-Archimedean potential associated with $\text{Tr}_E(\Gamma)$ is not very easy to describe. We will only do this in the following special but important case.

THEOREM 4.21. *Assume that X is smooth projective and $\{\theta\}$ is the first Chern class of a pseudoeffective line bundle L and there is a smooth Hermitian metric h_0 with $c_1(L, h_0) = \theta$. Consider an element $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ with non-trivial restriction to E . Then*

$$(4.7) \quad \text{Tr}_E(\Gamma)^{\text{an}}|_{E^{\text{div}}} = \Gamma^{\text{an}}|_{E^{\text{div}}}.$$

Observe that there is a canonical identification $E^{\text{div}} = \tilde{E}^{\text{div}}$.

In order to prove the theorem, we need to recall the following notion introduced in [DXZ23].

Definition 4.22. A *piecewise linear curve* ψ_\bullet in $\text{PSH}(X, \theta)$ associated with $\psi_{\tau_j} \in \text{PSH}(X, \theta)$, for a finite number of parameters $\tau_0 > \tau_1 > \dots > \tau_N$ is the affine interpolation of this data:

- (1) $\psi_\tau = \psi_{\tau_N}$ for $\tau \leq \tau_N$;
- (2) For $t \in (0, 1)$ and $i = 0, \dots, N-1$, we have $\psi_{(1-t)\tau_i + t\tau_{i+1}} = (1-t)\psi_{\tau_i} + t\psi_{\tau_{i+1}}$;
- (3) $\psi_\tau = -\infty$ for $\tau > \tau_0$.

Similarly, an *essentially piecewise linear curve* in $\text{PSH}(X, \theta)$ is a curve η_\bullet in $\text{PSH}(X, \theta) \cup \{-\infty\}$ parameterized by \mathbb{R} such that there is a piecewise linear curve ψ_\bullet in $\text{PSH}(X, \theta)$ as above such that $\eta_\tau \sim_{\mathcal{I}} \psi_\tau$ for all $\tau \in \mathbb{R}$.

The analytification ψ^{an} of a piecewise linear or essentially piecewise linear curve ψ_\bullet as above is defined as

$$(4.8) \quad \psi^{\text{an}}(v) := \sup_{\tau \leq \tau_0} (\psi_\tau^{\text{an}}(v) + \tau) = \max_{\tau_i} (\psi_{\tau_i}^{\text{an}}(v) + \tau_i) \quad \text{for } v \in X^{\text{an}}.$$

PROOF OF THEOREM 4.21. We may assume that $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$. Let $\phi = \Gamma^{\text{an}} \in \text{PSH}(L^{\text{an}})$. By Lemma 4.19, $\phi(v_{E, \text{triv}}) \neq -\infty$.

Take an ample line bundle S on X , a Kähler form ω in $c_1(S)$. Write ϕ as the decreasing limit of a sequence ϕ^i of elements in $\mathcal{H}_{\mathbb{Q}}^{\text{gf}}(L + i^{-1}S)$. This is possible by [BJ21, Corollary 12.18].

Take $\Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta + i^{-1}\omega)$ such that $\Gamma^{i, \text{an}} = \phi^i$. Note that by Lemma 3.10, $\Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta + i^{-1}\omega)_{>0}$.

It follows from Lemma 4.10 (applied to the images of Γ^i in $\text{PSH}^{\text{NA}}(X, \theta + \omega)$) that for any $\tau < \Gamma_{\max}$, we have

$$\lim_{i \rightarrow \infty} \Gamma_\tau^i = \Gamma_\tau.$$

In particular, by Lemma 2.10, the Lelong numbers of Γ_τ^i converge to those of Γ_τ .

It follows that $\Gamma_\tau^i \xrightarrow{d_{S, \theta + \omega}} \Gamma_\tau$ for all $\tau < \Gamma_{\max}$.

By Lemma 4.19 again, each Γ^i has non-trivial restriction to E . By [DX24, Lemma 3.12], we have

$$\text{Tr}_E(\Gamma^i)_\tau^{(\theta + \omega)|_{\tilde{E}}} \xrightarrow{d_S} \text{Tr}_E(\Gamma)_\tau^{(\theta + \omega)|_{\tilde{E}}}$$

for any $\tau < (\text{Tr}_E(\Gamma))_{\max}$. Thanks to [Xia22, Theorem 0.10], the Lelong numbers of $\text{Tr}_E(\Gamma^i)_\tau^{(\theta + \omega)|_{\tilde{E}}}$ converge to those of $\text{Tr}_E(\Gamma)_\tau^{(\theta + \omega)|_{\tilde{E}}}$. Therefore, by [DXZ23, Lemma 3.13],

$$\text{Tr}_E(\Gamma)^{\text{an}}(c \text{ord}_F) = \inf_i \text{Tr}_E(\Gamma^i)^{\text{an}}(c \text{ord}_F)$$

for any $c \text{ord}_F \in E^{\text{div}}$. In particular, it suffices to prove (4.7) with Γ^i in place of Γ .

In other words, we have reduced to the case where $\phi \in \mathcal{H}_{\mathbb{Q}}^{\text{gf}}(L)$ and L is big.

Under this extra assumption, as proved in the proof of [DXZ23, Theorem 3.12], there is a piecewise linear curve ψ connecting potentials lying in the space of potentials generated by functions of the form

$$(4.9) \quad m^{-1} \log \max_i (|s_i|_{h_0}^2 + c)$$

using convex combinations and finite maxima, where $m \in \mathbb{Z}_{>0}$, the s_i 's are finitely many holomorphic sections of $L^{\otimes m}$ and $c \in \mathbb{Q}$ such that Γ_{τ} is the \mathcal{I} -envelope of η_{τ} for any $\tau < \Gamma_{\max}$, where η is the concave envelope of ψ . Moreover, after adjusting ψ , we may guarantee that ψ itself is concave. In particular, $\Gamma_{\tau} \sim_{\mathcal{I}} \psi_{\tau}$ for all $\tau < \Gamma_{\max}$.

It follows that for any $c \text{ord}_F \in E^{\text{div}}$,

$$\begin{aligned} \phi|_{E^{\text{an}}}(c \text{ord}_F) &= \sup_{\tau < \Gamma_{\max}} (\psi_{\tau}^{\text{an}}(c \text{ord}_F) + \tau) \\ &= \sup_{\tau < \Gamma_{\max}} ((\psi_{\tau}|_{\bar{E}})^{\text{an}}(c \text{ord}_F) + \tau) \\ &= \sup_{\tau < \Gamma_{\max}} (\text{Tr}_E(\psi_{\tau})^{\text{an}}(c \text{ord}_F) + \tau) \\ &= \sup_{\tau < \Gamma_{\max}} (\text{Tr}_E(\Gamma_{\tau})^{\text{an}}(c \text{ord}_F) + \tau) \\ &= \text{Tr}_E(\Gamma)^{\text{an}}(c \text{ord}_F). \end{aligned}$$

Here we adopt the convention that $(-\infty)^{\text{an}}(c \text{ord}_F) = -\infty$ to simplify the notations. The third equality follows from the linearity of the trace operator [DX24, Lemma 3.10]. The fourth equality follows from [DX24, Lemma 3.9]. In order to justify the second equality, it suffices to show that for $\tau < (\text{Tr}_E(\Gamma))_{\max}$, if ψ_{τ} has the form (4.9), then

$$(\psi_{\tau}|_{\bar{E}})^{\text{an}}(\text{ord}_F) = \psi_{\tau}^{\text{an}}(\text{ord}_F).$$

We could easily reduce to the case $m = 1$ and $\psi_{\tau} = \log |s|_{h_0}^2$ for a single holomorphic section s of L which does not vanish identically on E . In this case, the above equality is trivial. \square

In particular, the trace operator is compatible with the partial order and the addition by a constant.

As an interesting corollary,

Corollary 4.23. *Assume the same assumptions as in Theorem 4.21. Suppose that $\phi \in \text{PSH}(L^{\text{an}})$ is a non-Archimedean metric represented by $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ and E is an irreducible reduced subvariety of X . Then*

$$\phi(v_{E, \text{triv}}) = \sup \{ \tau \in \mathbb{R} : \nu(\Gamma_{\tau}^{\theta+\omega}, E) = 0 \}$$

for any Kähler form ω on X .

4.8. Functoriality. Let $\pi: Y \rightarrow X$ be a proper bimeromorphic morphism from a reduced irreducible unibranch compact Kähler space of dimension n . By Zariski's main theorem, the obvious pull-back map induces a bijection

$$\pi^*: \text{PSH}(X, \theta) \xrightarrow{\sim} \text{PSH}(Y, \pi^*\theta).$$

In particular, it induces a canonical bijection

$$\pi^*: \text{PSH}^{\text{NA}}(X, \theta)_{>0} \xrightarrow{\sim} \text{PSH}^{\text{NA}}(Y, \pi^*\theta)_{>0}$$

as follows: the image of $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ is given by

$$\pi^*\Gamma: (-\infty, \Gamma_{\max}) \rightarrow \text{PSH}(Y, \pi^*\theta), \quad (\pi^*\Gamma)_\tau := \pi^*\Gamma_\tau.$$

This induces a canonical bijection

$$\pi^*: \text{PSH}^{\text{NA}}(X, \theta) \xrightarrow{\sim} \text{PSH}^{\text{NA}}(Y, \pi^*\theta).$$

The injectivity is trivial and the surjectivity follows from Lemma 3.8.

Note that there is a canonical identification $Y^{\text{div}} \xrightarrow{\sim} X^{\text{div}}$, so that we can identify $(\pi^*\Gamma)^{\text{an}}$ and Γ^{an} .

One trivially verifies that all the previous operations are functorial with respect to π^* .

5. Comparison with Boucksom–Jonsson’s theory

Let X be a reduced irreducible unibranch projective variety and L be a pseudoeffective line bundle on X with $\xi = c_1(L)$. Fix a smooth Hermitian metric h_0 on L with $\theta = c_1(L, h_0)$.

Recall the following basic result:

THEOREM 5.1 ([BJ21, Theorem 4.22]). *Let $\phi, \psi \in \text{PSH}(L^{\text{an}})$. Assume that $\phi \leq \psi$ on X^{div} , then the same holds on X^{an} .*

In particular, as we have computed the effect of our operations on X^{div} , their behaviours on X^{an} are in fact completely determined.

Corollary 5.2. *The order relation, addition by a constant, maximum, decreasing limit, rescaling and addition correspond to the corresponding operations in Boucksom–Jonsson’s theory. Assume that the envelope conjecture holds for X , then the same holds for the supremum operation.*

Briefly speaking, we have established the correspondence between the non-Archimedean pluripotential theory and the convex geometry of complex potentials as in Table 1.

Finally, we have the following comparison result between our theory of non-Archimedean potentials and Boucksom–Jonsson’s.

Corollary 5.3. *The following are equivalent:*

- (1) *The envelope conjecture of Boucksom–Jonsson holds for (X, L) ;*
- (2) *The comparison map $\Gamma \mapsto \Gamma^{\text{an}}$ defined in Definition 3.11 admits an extension as a bijection $\text{PSH}^{\text{NA}}(X, \theta) \xrightarrow{\sim} \text{PSH}(L^{\text{an}})$.*

PROOF. By Theorem 5.1, if the map in (2) exists, it is necessarily unique.

Let $\pi: Y \rightarrow X$ be a projective resolution. Consider the following diagram:

$$\begin{array}{ccc} \text{PSH}^{\text{NA}}(X, \theta) & \overset{\text{an}}{\cdots\cdots\cdots} & \text{PSH}(L^{\text{an}}) \\ \downarrow \pi^* & & \downarrow \pi^* \\ \text{PSH}^{\text{NA}}(Y, \pi^*\theta) & \xrightarrow{\text{an}} & \text{PSH}(\pi^*L^{\text{an}}). \end{array}$$

Clearly, (2) means that we can complete the dotted map as a bijection so that the diagram is commutative. We already know that the left-vertical map is a bijection, see Section 4.8. The lower-horizontal map is also a bijection by Theorem 3.12.

Now assume (1), then the right-vertical map is also a bijection by [BJ21, Lemma 5.13]. Therefore, (2) follows.

Conversely, assume that (2) holds. Take an increasing net $\phi_i \in \text{PSH}^{\text{NA}}(L^{\text{an}})$ with $\phi_i \leq 0$. Recall that the envelope conjecture means that ϕ_i converges to some $\phi \in \text{PSH}^{\text{NA}}(L^{\text{an}})$ with respect to the pointwise convergence on X^{div} . We write $\phi_i = \Gamma^{i, \text{an}}$ for some $\Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta)$. It suffices to take $\phi = (\sup^*_i \Gamma^i)^{\text{an}}$ and apply Theorem 4.13. \square

References

- [BBJ21] Robert J. Berman, Sébastien Boucksom, and Mattias Jonsson, *A variational approach to the Yau-Tian-Donaldson conjecture*, J. Amer. Math. Soc. **34** (2021), no. 3, 605–652. MR 4334189
- [Ber93] Vladimir G. Berkovich, *Étale cohomology for non-Archimedean analytic spaces*, Inst. Hautes Études Sci. Publ. Math. (1993), no. 78, 5–161 (1994). MR 1259429
- [BFJ15] Sébastien Boucksom, Charles Favre, and Mattias Jonsson, *Solution to a non-Archimedean Monge-Ampère equation*, J. Amer. Math. Soc. **28** (2015), no. 3, 617–667. MR 3327532
- [BFJ16] ———, *Singular semipositive metrics in non-Archimedean geometry*, J. Algebraic Geom. **25** (2016), no. 1, 77–139. MR 3419957
- [BHH17] Sébastien Boucksom, Tomoyuki Hisamoto, and Mattias Jonsson, *Uniform K-stability, Duistermaat-Heckman measures and singularities of pairs*, Ann. Inst. Fourier (Grenoble) **67** (2017), no. 2, 743–841. MR 3669511
- [BJ21] Sébastien Boucksom and Mattias Jonsson, *Global pluripotential theory over a trivially valued field*, 2021.
- [BJ22] ———, *Addendum to the article global pluripotential theory over a trivially valued field*, 2022.
- [CL06] Antoine Chambert-Loir, *Mesures et équidistribution sur les espaces de Berkovich*, J. Reine Angew. Math. **595** (2006), 215–235. MR 2244803
- [CLD12] Antoine Chambert-Loir and Antoine Ducros, *Formes différentielles réelles et courants sur les espaces de Berkovich*, 2012.
- [DDNL18] Tamás Darvas, Eleonora Di Nezza, and Chinh H. Lu, *Monotonicity of nonpluripolar products and complex Monge-Ampère equations with prescribed singularity*, Anal. PDE **11** (2018), no. 8, 2049–2087. MR 3812864
- [DDNL21] Tamás Darvas, Eleonora Di Nezza, and Hoang-Chinh Lu, *The metric geometry of singularity types*, J. Reine Angew. Math. **771** (2021), 137–170. MR 4234098
- [Dem85] Jean-Pierre Demailly, *Mesures de Monge-Ampère et caractérisation géométrique des variétés algébriques affines*, Mém. Soc. Math. France (N.S.) (1985), no. 19, 124. MR 813252
- [DX21] Tamás Darvas and Mingchen Xia, *The volume of pseudoeffective line bundles and partial equilibrium*, Geom. Topol. (2021).
- [DX22] Tamás Darvas and Mingchen Xia, *The closures of test configurations and algebraic singularity types*, Adv. Math. **397** (2022), Paper No. 108198, 56. MR 4366232
- [DX24] Tamás Darvas and Mingchen Xia, *The trace operator of quasi-plurisubharmonic functions on compact Kähler manifolds (to appear)*, 2024.
- [DXZ23] Tamás Darvas, Mingchen Xia, and Kewei Zhang, *A transcendental approach to non-Archimedean metrics of pseudoeffective classes*, 2023.
- [GK20] Patrick Graf and Tim Kirschner, *Finite quotients of three-dimensional complex tori*, Ann. Inst. Fourier (Grenoble) **70** (2020), no. 2, 881–914. MR 4105953
- [Gub16] Walter Gubler, *Forms and current on the analytification of an algebraic variety (after Chambert-Loir and Ducros)*, Nonarchimedean and tropical geometry, Simons Symp., Springer, [Cham], 2016, pp. 1–30. MR 3700066
- [Li20] Chi Li, *Geodesic rays and stability in the cscK problem*, 2020.
- [Roc70] R. Tyrrell Rockafellar, *Convex analysis*, Princeton Mathematical Series, No. 28, Princeton University Press, Princeton, N.J., 1970. MR 0274683
- [RWN14] Julius Ross and David Witt Nyström, *Analytic test configurations and geodesic rays*, J. Symplectic Geom. **12** (2014), no. 1, 125–169. MR 3194078
- [SD18] Zakarias Sjöström Dyrefelt, *K-semistability of cscK manifolds with transcendental cohomology class*, J. Geom. Anal. **28** (2018), no. 4, 2927–2960. MR 3881961

- [Thu05] Amaury Thuillier, *Théorie du potentiel sur les courbes en géométrie analytique non archimédienne. applications à la théorie d'arakov*, Ph.D. thesis, Université Rennes 1, 2005.
- [WN19] David Witt Nyström, *Monotonicity of non-pluripolar Monge-Ampère masses*, Indiana Univ. Math. J. **68** (2019), no. 2, 579–591. MR 3951074
- [Xia21a] Mingchen Xia, *Mabuchi geometry of big cohomology classes with prescribed singularities*, 2021.
- [Xia21b] ———, *Partial Okounkov bodies and Duistermaat–Heckman measures of non-Archimedean metrics*, 2021.
- [Xia22] ———, *Non-pluripolar products on vector bundles and Chern–Weil formulae*, Math. Ann. (2022).
- [Xia23a] ———, *Mabuchi geometry of big cohomology classes*, J. Reine Angew. Math. **798** (2023), 261–292. MR 4579706
- [Xia23b] ———, *Pluripotential-theoretic stability thresholds*, Int. Math. Res. Not. IMRN (2023), no. 14, 12324–12382. MR 4615232
- [Zha95] Shouwu Zhang, *Positive line bundles on arithmetic varieties*, J. Amer. Math. Soc. **8** (1995), no. 1, 187–221. MR 1254133

INSTITUT DE MATHÉMATIQUES DE JUSSIEU – PARIS RIVE GAUCHE, 4 PLACE JUSSIEU, PARIS
Email address: mingchen@imj-prg.fr